

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 336 200**  
**A1**

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 89105004.9

(51) Int. Cl.4: A61K 31/54 , A61K 9/06

(22) Date of filing: 21.03.89

Claims for the following Contracting States: ES  
+ GR.

(30) Priority: 01.04.88 IT 2007388

(43) Date of publication of application:  
11.10.89 Bulletin 89/41

(84) Designated Contracting States:  
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: CHIESI FARMACEUTICI S.p.A.  
Via Palermo, 26/A  
I-43100 Parma(IT)

(72) Inventor: Chiesi, Paolo  
Via Palermo, 26/A  
I-43100 Parma(IT)  
Inventor: Pavesi, Luciana  
Via Palermo, 26/A  
I-43100 Parma(IT)

(74) Representative: Minoja, Fabrizio  
Studio Consulenza Brevettuale Via Rossini,  
8  
I-20122 Milano(IT)

(54) Pharmaceutical compositions of piroxicam in aqueous solutions and process for their preparation.

(57) Aqueous pharmaceutical formulations containing the active principle piroxicam or N-(2-pyridyl)-2-methyl-4-hydroxy-2H-1,2-benzothiazine-3-carboxyamido-1,1-dioxide, are described.

The compositions of the invention are particularly suited for the injectable use, both because of their tolerability degree and for the favourable pharmacokinetics profile and may be advantageously used also for the preparation of collyria or eye-washes.

EP 0 336 200 A1

# PHARMACEUTICAL COMPOSITIONS OF PIROXICAM IN AQUEOUS SOLUTIONS AND PROCESS FOR THEIR PREPARATION

The present invention refers to pharmaceutical compositions of piroxicam in aqueous solutions, particularly suited for the preparation of injectable compositions.

Piroxicam, whose chemical name is N-(2-pyridyl)-2-methyl-4-hydroxy-2H-1,2-benzothiazine-3-carboxyamido-1,1-dioxide, is a nonsteroidal antiinflammatory drug widely used in the treatment of arthrorheumatic diseases; for these indications, it is normally administered by the oral or rectal route.

Piroxicam is, on the other side, endowed with a remarkable analgesic activity and therefore it is effectively used also in the treatment of pain both associated with arthrorheumatic diseases and of different kind (cephalea, dysmenorrhea, painful manifestations of different kind).

When the desired therapeutic effect is mainly the analgesic one, it is particularly useful to make available compositions for parenteral use allowing more rapid hematic levels of active principle and consequently a rapid alleviation of pain. The poor water-solubility of piroxicam does not allow the formulation thereof in aqueous solutions suited for the preparation of injectable compositions.

In order to cope with this drawback, several proposals have been made.

The European Patent 66458 discloses piroxicam salts with arginine and lysine, which may be used for the preparation of compositions for parenteral use.

Analogously, the European Patent 66459 discloses piroxicam salts with ethylenediamine, monoethanolamine and diethanolamine.

EP-A-177870 discloses aqueous-organic solutions of piroxicam, consisting of 50% water, 40% ethanol to which sub-stoichiometric amounts of N-methylglucamine are added as stabilizer of the solution.

It has now been found a piroxicam composition for parenteral use wherein the active principle is present as alkaline or alkali-earth metal salt and preferably as sodium salt, dissolved in water for injectable preparations.

This formulation, which may also be used for the preparation of collyria or eye-washes, besides to the preparation of injectable solutions for intramuscular, intravenous or intraarticular use, has the following advantages in comparison with the known compositions:

1. easy preparation, because the process allows to obtain the final formulation by few and simple steps, without the isolation and crystallization of salts of the active principle;

2. total absence of cosolvents such as alcohols or glycols which may give rise to general or local tolerability problems;

3. as a consequence of what above said in item 2, a better respect of the physiological conditions and consequently a particularly high tolerability;

4. a better pharmacokinetic behaviour, mainly as far as the maximal values of plasmatic concentrations ( $C_{max}$ ) are concerned, in comparison to a commercial piroxicam injectable preparation.

A second object of the invention is provided by the process for the preparation of aqueous solutions of piroxicam, consisting in the dispersion in water of the active principle in concentrations ranging from 1 to 4% w/v and subsequent addition, under stirring, of an aqueous solution of alkaline or alkali-earth hydroxide, preferably 1N sodium hydroxide in stoichiometric amount up to complete dissolution.

Glycine is added to said solution in amounts ranging from 8-10% w/v as pH-stabilizer.

The so obtained final solution may be subdivided in vials and subsequently sterilized in autoclave or alternatively it may be sterilized by filtration and then dosed in vials and lyophilized in sterile room. The lyophilized may be easily re-dissolved in water for injectable preparations, just prior to the use, giving a clear aqueous solution.

The latter preparation is preferred since the active principle, in form of lyophilized powder, keeps its chemical stability unchanged during time.

The following examples illustrate but do not limit the invention anyhow.

## **EXAMPLE 1: Preparation of the lyophilized**

Composition of a lyophilized vial:

Piroxicam	mg 20
Sodium hydroxide 1N	ml 0,12
Glycine	mg 100
Water for injection	q.s. to 1 ml

#### Preparation method

Piroxicam is dispersed in water, the aqueous solution of sodium hydroxide is added so as to obtain a clear solution.

The glycine is added and the solution is stirred till it becomes clear.

The solution is then sterile-filtered through a 0.2  $\mu$ m membrane.

The subsequent operations of distribution, lyophilization and vial sealing are carried out in sterile room by known and standardized methods.

The lyophilized, prior to its use, is dissolved in 2 ml of water for injection yielding a clear solution having pH = 8-8.5.

#### EXAMPLE 2: Preparation of the ready-to-use solution

Injectable piroxicam compositions as ready-to-use solutions are prepared as in Example 1.

The solution may conveniently contain benzyl alcohol as a conservative and local anesthetic agent.

Two different composition are reported:

Composition a)	
Piroxicam	mg 20
Sodium hydroxide 1N	ml 0.172
Glycine	mg 80
Water for injection	q.s. to 2 ml
Composition b)	
Piroxicam	mg 20
Sodium hydroxide 1N	ml 0.172
Glycine	mg 80
Benzyl alcohol	mg 20
Water for injection	q.s. to 2 ml

The sterilization may be carried out by filtration through a 0.2  $\mu$ m membrane and subsequent distribution in vials in sterile room. Alternatively, the sterilization may be carried out by heating in autoclave after distribution of the solution in vials.

Said aqueous solutions of piroxicam may also be used in the preparation of collyria, optionally adding to the composition antibacterial compounds, such as quaternary ammonium salts.

The kinetic comparison of the pharmaceutical composition of the invention has been evaluated in man in comparison with a standard, commercially available injectable formulations whose composition is the following:

Piroxicam	mg 20
Benzyl alcohol	mg 20
Nicotinamide	mg 30
Propylene glycol	mg 400
Water for injection	ml 1

The study was carried out on 6 adult healthy volunteers, according to a cross-over design using the injectable composition of Example 1, hereinafter referred to as CHF 1251.

Both the reference composition and that of the present invention, comprised 20 mg of active principle. The administration was carried out by the intramuscular route. The determination of the plasma concentration of the active principle at different times from administration was carried out by high pressure liquid chromatography (HPLC).

The results obtained expressed as mean values of the plasma concentrations at different times, AUC (area under the curve blood concentration/time), MRT (mean residence time),  $C_{max}$  (maximum plasma concentration) and  $T_{max}$  (time needed for reaching the maximum plasma concentration) are reported in table 1.

5

Composi- tions	No. subjects	Meanvalue and standard deviating ( $\bar{X}$ , $\pm$ SD) of plasma conc. of ( $\mu\text{g/ml}$ ) of piroxicam at different times (h) after administration									AUC	MRT	$C_{\text{max}}$	$T_{\text{max}}$
		0.25	0.50	1	2	4	8	24	h	$\mu\text{g/ml}$				
CHF 1251	6	$\bar{X}$	2.38	2.43	2.43	2.02	1.73	1.79	1.26	115.28	60.20	2.61	1.58	
		$\pm$ D.S.	0.23	0.48	0.51	0.37	0.42	0.46	0.41	29.02	11.29	0.39	2.18	
Commercial injectable piroxicam	6	$\bar{X}$	0.98	1.57	1.89	1.69	1.92	1.78	1.35	150.63	54.29	2.20	3.58	
		$\pm$ D.S.	0.32	0.52	0.34	0.30	0.53	0.54	0.44	71.35	29.52	0.33	2.38	

The AUC and Cmax values were compared by the variance analysis method as disclosed by Wagner J.G. on *Fundamental of Clinical Pharmacokinetics*, Hamilton 3<sup>rd</sup>, Drug Intell. Publ. 291-4, 1975.

The Tmax values were composed by the non-parametric Wilcoxon test as disclosed by Siegel S. on "Statistica non parametrica per le scienze del comportamento", Ed. OS, Firenze 63-70, 1966.

As it can be shown from the data reported in Table 1, the absorption is very fast after administration of the injectable piroxicam formulation CHF 1251: the blood concentration peak is ranging from 2.14 to 3.14 µg/ml (mean value 2.61 µg/ml) and it is achieved after a mean time of 1.58 h after treatment whereas the commercial composition, whose mean max. concentration was 2.20 µg/ml achieved after a mean time of 3.58 h after treatment.

The plasmatic profile of CHF 1251 shows moreover that the drug remains in the plasma compartment for a long time, as it is evident from MRT values ranging from 43 to 74 hours (mean value: 60 hours and 20 minutes).

The permanence time of piroxicam in plasma after CHF 1251 administration is longer than that obtained with the commercial injectable formulation of piroxicam, showing a mean value of about 54 hours and 30 minutes.

According to the results obtained in this comparative study, using as a reference drug a commercial, standard injectable preparation of piroxicam, it can be concluded that the composition of the invention has a particularly favourable plasma pharmacokinetics profile, characterized by a fast absorption and above all by the achievement of a significantly higher maximum plasma concentration (about 5% higher) than that obtained with the reference formulation.

This result is particularly important for the desired therapeutic effect of a fast and effective analgesic action.

In fact, the treatment by parenteral route is used in cases of acute pain of remarkable seriousness just for the faster rapidity of the antalgic effect.

In the case of non-steroidal antiphlogistic drugs, clinically used for the treatment of acute pain of different etiology, the parenteral administration is generally better tolerated in comparison with the oral route as far as undesired side-effects at the gastro-intestinal tract level are concerned and it is preferably used for patients affected by gastro-intestinal diseases.

In addition to a better general tolerability in comparison to the oral route, the composition of the invention has also a better local tolerability in comparison with the commercial, standard injectable formulation, already used for the kinetics study.

Most of the non-steroidal anti-inflammatory drugs usually administered by intramuscular route as well as other active principles are badly tolerated locally since they are endowed with an histolesive action causing the release of neurohumors and of intracellular proteolytic enzymes, with consequent local intolerance manifestations such as burning, pain, reddening and induration at the injection site.

One of the quantitative evaluation methods of the tissue damage is the determination of the serum level of an enzyme characteristics of the muscular fibre, the creatine phosphokinase (CPK), normally present in blood in low concentrations (up to 250 mU/ml), which may however considerably increase with few hours in case of myolysis and cellular necrosis due to causes of different nature such as the administration of drugs by the intramuscular route.

The administration of serum level of CPK represents therefore a convenient biochemical test whenever the local tolerability of any injectable preparation for intramuscular use should be evaluated (Cingolani E. et al. *Il Farmaco* Ed. Pr. 41(3), 89, 1986).

The evaluation of the local tolerability of the composition of the invention has been carried out by measuring the serum levels of CPK and the direct detection of intolerance symptoms and signs at the injection site, in 6 healthy volunteers which were considered suited according to a series of exam and were subjected to a treatment cycle with 6 vials of CHF 1251 (2 vials at 8 hours interval at the first day and 1 vial/day for the subsequent 4 days).

The tolerability of the reference drug, i.e. of the commercial injectable formulation of piroxicam was evaluated by the same method and at the same time.

The results of the serum CPK levels at different times are shown in Table 2.

**TABLE 2** Meanvalues and standard deviation of plasma CPK (mU/ml) at different times during a treatment cycle respectively of CHF 1251 and of commercial injectable formulation of piroxicam.

TIMES														
DRUG	No. subjects	0	1	2	3	4	5	6	7	8	9	10	11	12
CHF 1251	6	81.2	77.0	79.5	95.7	98.8	100.5	94.0	98.7	103.3	114.2	106.8	93.7	81.3
	D.S.	6.0	20.4	19.7	18.5	21.4	34.7	12.4	27.5	22.8	27.0	18.8	19.2	12.6
commercial injectable piroxicam	6	135.3	140.0	168.7	191.5	306.3	334.5	334.5	196.3	201.8	220.2	234.5	242.2	198.5
	D.S.	47.9	47.0	40.4	59.7	140.1	164.3	164.3	73.0	71.1	99.2	110.2	103.4	89.8

Before test start: 0 = basal

During test:

1 = day 1, time 0  
 2 = day 1, time 20'  
 3 = day 1, time 7 hours  
 4 = day 1 (second injection), time 2 h  
 5 = day 3, time 20'  
 6 = day 3, time 7 hours  
 7 = day 5, time 0  
 8 = day 5, time 20'  
 9 = day 5, time 2 hours  
 10 = day 5, time 3 hours  
 11 = day 5, time 7 hours  
 12 = day 6, time 0

As it can be noticed from the data of table 2, the histolesive effect is completely lacking with CHF 1251 whereas it is present, although in modest amount, with the reference formulation.

Since the two formulations comprise the same active principle, said effect is probably due to the solvent's action (a mixture of propylene glycol-benzyl alcohol-water) of the commercial product.

CHF 1251 using a purely aqueous carrier did not induce any tissue damage detectable by the used tests.

This objective evaluation of the local tolerability of the composition of the invention has been further confirmed by the subjective evaluation of volunteers who only in some cases complained about burning reactions at the injection site.

## Claims

1. Pharmaceutical compositions of piroxicam in aqueous solution for injectable or ophthalmic use, wherein piroxicam is present as alkaline or alkali earth-metal salt.

2. Pharmaceutical compositions according to claim 1, for injectable use, containing from 10 to 30 mg of active principle per dosage unit, in form of ready-to-use solution or in lyophilized form to be dissolved in water for injection before use.

3. Pharmaceutical compositions according to claim 1, for ophthalmic use, wherein the concentration of the active principle is from 0.2 to 1%.

4. A process for the preparation of pharmaceutical compositions of claims 1-3 comprising:

a) dispersing the active principle in water in concentrations from 1 to 4% w/v;

b) adding under stirring an aqueous solution of an alkaline or alkaline-earth metal hydroxide in stoichiometric amount up to complete dissolution;

c) adding glycine in amounts ranging from 8 and 10% w/v, as pH stabilizer.

Claims for the following Contracting States: GR, ES

1. A process for the preparation of piroxicam in aqueous solution for injectable or ophthalmic use, wherein piroxicam is present as alkaline or alkali earth-metal salt, comprising:

a) dispersing the active principle in water in concentrations from 1 to 4% w/v;

b) adding under stirring an aqueous solution of an alkaline or alkaline-earth metal hydroxide in stoichiometric amount up to complete dissolution;

c) adding glycine in amounts ranging from 8 and 10% w/v, as pH stabilizer.



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number

EP 89 10 5004

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Y	CH-A- 594 650 (PFIZER) * Claims * ---	1-4	A 61 K 31/54 A 61 K 9/06
Y	CHEMICAL ABSTRACTS, vol. 99, no. 12, 19th September 1983, page 338, abstract no. 93582y, Columbus, Ohio, US; C.D. HERZFELDT et al.: "Dissociation constant, solubilities and dissolution rates of some selected nonsteroidal antiinflammatories", & DRUG DEV. IND. PHARM. 1983, 9(5), 767-93 * Abstract * ---	1-4	
Y	CHEMICAL ABSTRACTS, vol. 107, no. 13, 28th September 1987, page 34, abstract no. 109008n, Columbus, Ohio, US; M. DE BERNARDI et al.: "Effects of topical treatment with piroxicam on prostaglandin synthesis in experimental uveitis", & INT. J. TISSUE REACT. 1987, 9(4), 361-4 * Abstract * -----	1-4	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			A 61 K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 26-06-1989	Examiner SCARPONI U.
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			